

THE MAXIMALITY OF THE CORE MODEL

E. SCHIMMERLING AND J.R. STEEL

1. INTRODUCTION

This paper deals with the core model, or K , as defined in [St1]. Our main results are interesting as part of the pure theory of this model, but they also yield some new consistency strength lower bounds, as we shall point out. Of course, one cannot give a sensible definition of K , much less prove anything about it, without an anti-large-cardinal hypothesis. For the most part, the hypothesis of this nature we shall assume here is that there is no inner model satisfying "There is a Woodin cardinal". The results we shall prove about K under this hypothesis have all been proved by others under more restrictive anti-large-cardinal hypotheses, and our proofs build on those earlier proofs.

The term "the core model" was introduced by Dodd and Jensen to mean, roughly, "the largest canonical inner model there is". They gave a precise definition of the term which captures this intuition in the case there is no inner model with a measurable cardinal in [DoJ]. Work of Mitchell, and then Martin, Mitchell and Steel, led to a precise definition which captures the intuition in the case there is no inner model with a Woodin cardinal, and every set has a sharp. (Cf. [St1]. The hypothesis that every set has a sharp seems to be a weakness in the basic theory of the core model for a Woodin cardinal. Theorem 5.1 shows how a weaker Erdős partition property suffices.) The canonicity of the core model is evidenced by the absoluteness of its definition: if G is set generic over V , then $K^V = K^{V[G]}$. (Cf. [St1, 7.3].) The maximality of the core model manifests itself in several ways. Perhaps the most important of these is the weak covering property of K , that it computes successors of singular cardinals correctly. (Cf. [St1, 8.15], [MiSchSt],

1991 *Mathematics Subject Classification.* 03E.

Key words and phrases. large cardinals, core models.

The research of the first author was partially supported by a National Science Foundation Mathematical Sciences Postdoctoral Research Fellowship at the Massachusetts Institute of Technology.

The research of the second author was partially supported by the National Science Foundation.

[MiSch], and [SchW].) Other manifestations of the maximality of K are its Σ_3^1 correctness (cf. [St1, 7.9]), and its rigidity (cf. [St1, 8.8]).

Our main result here is that K is maximal in another sense, namely, that every extender which "could be added" to K is already in K . More precisely, Theorem 2.3 says that K is maximal in the sense that any countably certified extender that coheres with K is actually on the extender sequence of K , and similarly for the iterates of K . Mitchell showed earlier that every countably complete K -ultrafilter is in K under the stronger hypothesis that there is no inner model with a measurable cardinal κ with $o(\kappa) = \kappa^{++}$. (Cf. [Mi]). It is not known if Theorem 2.3 holds for countably closed extenders. Theorem 2.5 is a result similar to Theorem 2.3 for mice with sufficiently small projectum.

Our first application of Theorem 2.3 is Theorem 3.1, which says that K computes successors of weakly compact cardinals correctly. This extends Kunen's result for L and the weak covering properties of K mentioned above.

Recall from [St1] that K is the transitive collapse of an elementary substructure of K^c , the background certified core model, and that K^c is the limit of the maximal, 1-small construction $\mathbb{C}$. In Theorem 3.2, another application of Theorem 2.3, we show that K^c and all the models on $\mathbb{C}$ are iterates of K .

Theorem 3.2 is used to prove Theorem 3.3, which says, roughly, that a Mahlo cardinal is a kind of closure point of the construction $\mathbb{C}$ (although not in as strong a sense as we would like).

The universality of initial segments of K is related to some open questions in descriptive set theory, such as the consistency strength of $u_2 = \aleph_2$. Jensen showed that it is consistent for a countable mouse to out-iterate $K \parallel \aleph_1$. Theorem 3.4, another application of maximality, shows that $K \parallel \kappa$ is universal for mice of height κ whenever κ is a cardinal $\geq \aleph_2$. (We proved Theorem 3.4 assuming countable closure, and Mitchell saw how to eliminate the assumption using [MiSch].)

Using Theorem 3.1 and [Sch2], we prove Theorem 4.1, which says that if κ is a weakly compact cardinal and $\square_\kappa^{<\omega}$ fails, then there is an inner model with a Woodin cardinal. The definition of $\square_\kappa^{<\omega}$ is recalled in Section 4; it is weaker than Jensen's principle $\square_\kappa$, and stronger than his $\square_\kappa^*$. Woodin noted that our proof, when combined with his core model induction, improves the conclusion of Theorem 4.1 to $L(\mathbb{R})$ -determinacy.

The proof of Theorem 4.1, together with the results in [Sch2] and [MiSchSt], are used in the proof of Theorem 4.2, where we show that if κ is a singular, strong limit cardinal, then there is an inner model with a Woodin cardinal. With little additional work, the conclusion

can be strengthened to the existence of an inner model with Woodin cardinals cofinal in its ordinals, which, by Woodin, implies Inductive Determinacy.

Some of the theorems in this paper have the hypothesis that Ω is a measurable cardinal. This is really just being used to see that the theory of [St1] applies (sharps for bounded subsets of Ω would do). For our theorems that have this hypothesis on Ω , Theorem 5.1 shows how to reduce it to the partition property: $\Omega \longrightarrow (\omega)_\alpha^{<\omega}$ for all $\alpha < \Omega$. For example, if it exists, the Erdős cardinal $\kappa(\omega)$ has this property, and it is consistent with $V = L$ that $\kappa(\omega)$ exists.

K is a particular model of the form $L[\vec{E}]$ where $\vec{E}$ is a coherent sequence of extenders. Section 6 pertains to models of the form $L[\vec{E}]$ in general. With no anti-large cardinal hypothesis, we show that if $L[\vec{E}]$ is adequately iterable, then it is also maximal in that $L[\vec{E}]$ satisfies the statement, "if F is a countably certified extender which coheres with $\vec{E}$, then $F = E_\alpha$ for some α ". One effect of this is that, typically, if $L[\vec{E}]$ satisfies a large cardinal property, then the large cardinal property is witnessed by extenders on $\vec{E}$.

2. MAXIMALITY

Definition 2.1. *Suppose that $\mathcal{M}$ is a premouse and that F is an extender that coheres with $\mathcal{M}$. Let $\kappa = \text{crit}(F)$ and $\nu = \nu(F)$. Suppose that*

$$\mathcal{A} \subseteq \bigcup_{n < \omega} \mathcal{P}([\kappa]^n)^{\mathcal{M}}.$$

*Then a **weak $\mathcal{A}$ -certificate for $(\mathcal{M}, F)$** is a pair (N, G) such that*

(a) N is a transitive, power admissible set, ${}^\omega N \subset N$,

$$\kappa \cup \mathcal{A} \cup \{\mathcal{J}_\kappa^{\mathcal{M}}\} \subset N,$$

and G is an extender over N ,

(b) $F \cap ([\nu]^{<\omega} \times \mathcal{A}) = G \cap ([\nu]^{<\omega} \times \mathcal{A})$,

(c) $V_{\nu+1} \subset \text{ult}(N, G)$,

(d) if $j : N \longrightarrow \text{ult}(N, G)$ is the ultrapower map, then $\mathcal{M}$ and $j(\mathcal{J}_\kappa^{\mathcal{M}})$ agree below $\text{lh}(F)$, and

(e) $\langle \mathcal{P}(\alpha) \cap N, \in \rangle \prec \langle \mathcal{P}(\alpha), \in \rangle$ whenever $\alpha < \kappa$.

Weak $\mathcal{A}$ -certificates are weaker than the $\mathcal{A}$ -certificates of [St1, 1.1] in two ways. First, we weakened the condition that $V_\kappa \subseteq N$ to just condition (e). And, second, $\text{ult}(N, G)$ is not required to be countably closed.

The following definition should be compared with [St1, 1.2].

Definition 2.2. Suppose that $\mathcal{M}$ is a premouse and F is an extender of length α that coheres with $\mathcal{M}$. Then $(\mathcal{M}, F)$ is **weakly countably certified** iff for all countable

$$\mathcal{A} \subseteq \bigcup_{n < \omega} \mathcal{P}([\kappa]^n) \cap J_\alpha^\mathcal{M}$$

there is a weak $\mathcal{A}$ -certificate for $(\mathcal{J}_\alpha^\mathcal{M}, F)$.

When $\mathcal{T}$ is an iteration tree, we shall sometimes write $\mathcal{M}(\mathcal{T}, \eta)$ for $\mathcal{M}_\eta^\mathcal{T}$, the η 'th model of $\mathcal{T}$, to avoid double superscripts. When $\mathcal{T}$ has a final model, we shall denote it either by $\mathcal{M}_\infty^\mathcal{T}$ or by $\mathcal{M}(\mathcal{T}, \infty)$. The following is our main maximality theorem.

Theorem 2.3. Suppose that Ω is a measurable cardinal and that there is no inner model with a Woodin cardinal. Let $\mathcal{T}$ be a normal iteration tree on K of successor length $< \Omega$. Suppose that F is an extender that coheres with $\mathcal{M}_\infty^\mathcal{T}$ and that

$$\text{lh}(E_\eta^\mathcal{T}) < \text{lh}(F)$$

for all $\eta < \text{lh}(\mathcal{T})$. Suppose that $(\mathcal{M}_\infty^\mathcal{T}, F)$ is weakly countably certified. Then F is on the $\mathcal{M}_\infty^\mathcal{T}$ -sequence.

The following notion shall enter into the proof of Theorem 2.3, and is of independent interest.

Definition 2.4. A premouse $\mathcal{M}$ is **conditionally α -strong** iff there is a normal iteration tree $\mathcal{T}$ on an α -strong premouse such that $\mathcal{T}$ has successor length, and there is an elementary embedding π of $\mathcal{M}$ into an initial segment of $\mathcal{M}_\infty^\mathcal{T}$ with $\text{crit}(\pi) \geq \alpha$.

Note that by a Löwenheim-Skolem argument, both $\mathcal{T}$ and $\mathcal{M}$ can be taken to have the same cardinality as $\mathcal{P}$ in Definition 2.4.

Proof of Theorem 2.3. Fix Ω , $\mathcal{T}$ and F as in the statement of Theorem 2.3 and let $\kappa = \text{crit}(F)$, $\nu = \nu(F)$, and $\lambda = \text{lh}(F)$.

Let μ_0 be an inaccessible cardinal $< \Omega$ such that $F \in V_{\mu_0}$. Let W be an A_0 -soundness witness for $\mathcal{J}_{\mu_0}^K$. Then $\mathcal{T}$ induces an iteration tree on W with the same extenders, drops, degrees, and tree structure as $\mathcal{T}$, and it is enough to show that F is on the sequence of the final model of the induced tree. For the rest of the proof, we shall write $\mathcal{T}$ for the induced iteration tree on W , and never again refer to the original iteration tree on K .

As in [St1, 9.7], we write $\Phi(\mathcal{T})$ for the phalanx derived from $\mathcal{T}$. The F -extension of $\Phi(\mathcal{T})$ is defined in [St1, 8.5]. By [St1, 8.6], we are done if we show that the F -extension of $\Phi(\mathcal{T})$ is $(\Omega + 1)$ -iterable. (Here we use

the hypothesis that $\mathcal{T}$ is normal.) For contradiction, suppose that $\mathcal{U}$ is an illbehaved iteration tree on the F -extension of $\Phi(\mathcal{T})$. We interpret $\mathcal{U} = \emptyset$ to mean that the ultrapower in the definition of the F -extension of $\Phi(\mathcal{T})$ is illfounded. By the tree property at Ω , $\text{lh}(\mathcal{U}) < \Omega$.

By the argument of [St1, 6.14], we may fix a successor cardinal μ such that $\mu_0 < \mu < \Omega$ and

- there is an iteration tree $\mathcal{S}$ on $\mathcal{J}_\mu^W$ with the same extenders, drops, degrees, and tree structure as $\mathcal{T}$, such that $\mathcal{M}_\eta^{\mathcal{S}}$ is an initial segment of $\mathcal{M}_\eta^{\mathcal{T}}$ whenever $\eta < \text{lh}(\mathcal{T})$, and
- there is an illbehaved iteration tree $\mathcal{V}$ on the F -extension of $\Phi(\mathcal{S})$ with the same extenders, drops, degrees, and tree structure as $\mathcal{U}$, such that $\mathcal{M}_\eta^{\mathcal{V}}$ is an initial segment of $\mathcal{M}_\eta^{\mathcal{U}}$ whenever $\eta < \text{lh}(\mathcal{U})$.

Recall that the F -extension of $\Phi(\mathcal{S})$ is the phalanx

$$\Phi(\mathcal{S}) \frown \langle \text{ult}_k(\mathcal{P}, F), k, \nu \rangle$$

where $\mathcal{P}$ is the initial segment of $\mathcal{M}_\gamma^{\mathcal{S}}$ to which F would be applied according to the rules for ω -maximal iteration trees, and $k < \omega$ is the degree. (This is not literally true, as the official definition of a phalanx, [St1, 9.5], has a fourth coordinate, in our case, either ν or λ , depending on the type of $\mathcal{P}$; we shall suppress this fourth coordinate.)

Let X be a countable elementary substructure of V_Ω with $\mathcal{V} \in X$ and let

$$\pi_X : M_X \longrightarrow V_\Omega$$

be the corresponding uncollapse. Say $\pi_X(\mathcal{S}_X) = \mathcal{S}$, $\pi_X(F_X) = F$, $\pi_X(\nu_X) = \nu$, $\pi_X(\lambda_X) = \lambda$, $\pi_X(\mathcal{P}_X) = \mathcal{P}$, and $\pi_X(\mathcal{V}_X) = \mathcal{V}$. By the absoluteness argument like that in the proof of [St1, 6.14], $\mathcal{V}_X$ is a simple, illbehaved iteration tree on

$$\Phi(\mathcal{S}_X) \frown \langle \text{ult}_k(\mathcal{P}_X, F_X), k, \nu_X \rangle,$$

the F_X -extension of $\Phi(\mathcal{S}_X)$.

Let Y be an elementary substructure of V_Ω with

$$X \cup (\lambda + 1) \subset Y$$

and $\text{card}(Y) = \text{card}(\nu)$. Let $\pi_Y : M_Y \longrightarrow V_\Omega$ be the uncollapse of Y and

$$\sigma = \pi_Y^{-1} \circ \pi_X.$$

Say $\pi_Y(\mathcal{S}_Y) = \mathcal{S}$ and $\pi_Y(\mathcal{P}_Y) = \mathcal{P}$. Note that $\mathcal{S}_Y$ has the same extenders, drops, degrees, and tree structure as $\mathcal{S}$, and that $\mathcal{M}(\mathcal{S}_Y, 0)$ is $\text{crit}(\pi_Y)$ -strong.

Let

$$\mathcal{A} = \text{ran}(\sigma) \cap \bigcup_{n < \omega} \mathcal{P}([\kappa]^n)^{\mathcal{P}_Y}$$

and (N, G) be a weak $\mathcal{A}$ -certificate for $(\mathcal{P}_Y, F)$. This is possible because $(\mathcal{M}_\infty^T, F)$ is weakly countably certified, and $\mathcal{P}_Y$ and $\mathcal{P}$ agree with $\mathcal{M}_\infty^T$ on subsets of κ . Let $j : N \longrightarrow \text{ult}(N, G)$ be the ultrapower map.

Pick a coordinate $b \in [\nu]^{<\omega}$ together with functions $u \mapsto \nu_u$, $u \mapsto \lambda_u$, and $u \mapsto \sigma_u$ in N such that

$$\begin{aligned}\nu &= [b, u \mapsto \nu_u]_G^N, \\ \lambda &= [b, u \mapsto \lambda_u]_G^N,\end{aligned}$$

and

$$\sigma = [b, u \mapsto \sigma_u]_G^N.$$

This is possible by clause (c) in Definition 2.1. By Łoś' theorem, there is a set $B \in G_b$ such that for all $u \in B$,

- $\sigma_u \upharpoonright |\mathcal{M}|$ is an elementary embedding of $\mathcal{M}$ into $\sigma_u(\mathcal{M})$ for every initial segment $\mathcal{M}$ of a model on $\Phi(\mathcal{S}_X)$,
- $\nu_u = \sigma_u(\nu_X) < \kappa$, and
- $\lambda_u = \sigma_u(\lambda_X) < \kappa$.

Lemma 2.4.1. *For G_b -almost every $u \in B$, there is a Σ_{k+1} -elementary, weak k -embedding*

$$\tau : \text{ult}_k(\mathcal{P}_X, F_X) \longrightarrow \mathcal{P}_Y$$

such that $\tau \upharpoonright \lambda_X = \sigma_u \upharpoonright \lambda_X$.

Proof. Let $\langle f_n \mid n < \omega \rangle$ be a sequence of $\Sigma_k(\mathcal{P}_Y)$ -functions and

$$\langle b_n \mid n < \omega \rangle$$

be a sequence of coordinates from $[\lambda]^{<\omega}$ such that

$$|\text{ult}_k(\mathcal{P}_Y, F)| \cap \text{ran}(\sigma) = \left\{ [b_n, f_n]_F^{\mathcal{P}_Y} \mid n < \omega \right\}$$

We assume that $b = b_0 \subset b_n \subset b_{n+1}$ for all $n < \omega$. We assume that f_n is chosen to be a constant function whenever possible. We assume that if

$$[b_n, f_n]_F^{\mathcal{P}_Y} = \sigma(\xi)$$

for some $\xi < \lambda_X$, then $\sigma(\xi) \in b_n$ and f_n is the projection function $u \mapsto u^{\sigma(\xi), \alpha}$.

For each $\xi < \lambda_X$ and $n \in \omega$ with $\sigma(\xi) \in b_n$, let

$$C_\xi^n = \left\{ u \in [\kappa]^{b_n} \mid \sigma_{u^{b_0, b_n}}(\xi) = u^{\sigma(\xi), b_n} \right\}.$$

Then $C_\xi^n \in G_{b_n}$. Since N is countably closed (or, just because the map $u \mapsto \sigma_u$ and all the reals are in N), it follows that

$$\langle C_\xi^n \mid \xi < \lambda_X \wedge \sigma(\xi) \in b_n \rangle$$

is an element of N .

If $n < \omega$ and e is the Gödel number for a Σ_k formula φ such that

$$\text{ult}_k(\mathcal{P}_Y, F) \models \varphi \left[[b_n, f_n]_F^{\mathcal{P}_Y} \right],$$

then let

$$T_e^n = \{u \in [\kappa]^{|b_n|} \mid \mathcal{P}_Y \models \varphi[f_n(u)]\}.$$

Then, $T_e^n \in F_{b_n} \cap \mathcal{A}$ by Łoś' theorem and our choice of $\mathcal{A}$. Moreover, the partial function $(n, e) \mapsto T_e^n$ is an element of N since N is countably closed.

Let $\langle B_n \mid n < \omega \rangle$ be a sequence in N such that $B_n \in G_{b_n}$ for all $n < \omega$, with the following three properties. First, $B_0 = B$. Second, for every $m, e < \omega$ such that T_e^m is defined, there is an $n \geq m$ with

$$(B_n)^{b_m, b_n} \subseteq T_e^m.$$

Third, for every $m < \omega$ and $\xi < \lambda_X$, there is an $n \geq m$ such that

$$(B_n)^{b_m, b_n} \subseteq C_\xi^m.$$

We need a “fiber” through the sets B_n . There is a tree $\mathbb{A}$ in N that searches for a function $g : \omega \longrightarrow [\kappa]^{<\omega}$ such that

$$g(m) \subset g(n) \in B_n$$

and

$$(b_n)^{g(m), g(n)} = b_m$$

whenever $m < n < \omega$. Since $\langle b_n \mid n < \omega \rangle$ is a branch through $j(\mathbb{A})_b$, there is a branch through $j(\mathbb{A})_b$ in $\text{ult}(N, G)$. Hence, for G_b -almost every $u \in B$, there is a branch g through $\mathbb{A}_u$ in N .

Given $u \in B$, fix $g \in N$ such that g is a branch through $\mathbb{A}_u$, and define a map

$$\tau : \text{ult}_k(\mathcal{P}_X, F_X) \longrightarrow \mathcal{P}_Y$$

by

$$\tau \left(\sigma^{-1} \left([b_n, f_n]_F^{\mathcal{P}_Y} \right) \right) = f_n(g(n)).$$

Because we “met” the sets T_e^n , τ is a weak k -embedding of $\text{ult}_k(\mathcal{P}_X, F_X)$ into $\mathcal{P}_Y$ (recall that weak k -embeddings need not be Σ_{k+1} -elementary). Because f_n was chosen to be the constant function whenever possible, if $i_X : \mathcal{P}_X \longrightarrow \text{ult}_k(\mathcal{P}_X, F_X)$ is the ultrapower map, then $\tau \circ i_X = \sigma$. Since i is a k -embedding and σ is elementary, τ is a Σ_{k+1} -elementary. Because we “met” the sets C_ξ^n , it follows that

$$\tau \upharpoonright \lambda_X = \sigma_u \upharpoonright \lambda_X.$$

□

Lemma 2.4.2. *There are functions $u \mapsto \mathcal{P}_u$ and $u \mapsto \tau_u$ in N such that for G_b -almost every $u \in B$, both V and N satisfy*

- (a) $\mathcal{P}_u$ is a conditionally $(\lambda_u + 1)$ -strong premouse,
- (b) $\mathcal{P}_u$ and $\mathcal{J}_\kappa^{\mathcal{P}}$ below $\lambda_u + 1$, and
- (c) τ_u is a Σ_{k+1} -elementary, weak k -embedding of

$$\text{ult}_k(\mathcal{P}_X, F_X)$$

into $\mathcal{P}_u$ such that $\tau_u \upharpoonright \lambda_X = \sigma_u \upharpoonright \lambda_X$.

Proof. First note that $\mathcal{P}_Y$ is conditionally $(\lambda + 1)$ -strong as witnessed by an initial segment of the iteration tree $\mathcal{S}_Y$ and the identity map on $\mathcal{P}_Y$.

Given $u \in B$ and a corresponding τ as in Lemma 2.4.1, let

$$\mathcal{P}_u^* = \mathcal{H}_\omega^{\mathcal{P}_Y}((\lambda_u + 1) \cup \text{ran}(\tau)),$$

and let

$$\tau_u^* : \text{ult}_k(\mathcal{P}_X, F_X) \longrightarrow \mathcal{P}_u^*$$

be the corresponding collapse of τ . Then the functions $u \mapsto \mathcal{P}_u^*$ and $u \mapsto \tau_u^*$ satisfy conditions (a), (b), and (c) in V , but these functions are not necessarily in N .

By the elementarity given in clause (e) of Definition 2.1, for each appropriate $u \in B$, there is a premouse $\mathcal{P}_u$ and an embedding τ_u , both in N , which satisfy conditions (a), (b), and (c) of Lemma 2.4.2 in both N and V . Since N satisfies AC, there are appropriate functions $u \mapsto \mathcal{P}_u$ and $u \mapsto \tau_u$ in N . $\square$

Fix functions as in Lemma 2.4.2 and let

$$\mathcal{P}^* = [b, u \mapsto \mathcal{P}_u]_G^N$$

and

$$\tau^* = [b, u \mapsto \tau_u]_G^N.$$

Lemma 2.4.3. τ^* is a Σ_{k+1} -elementary, weak k -embedding of

$$\text{ult}_k(\mathcal{P}_X, F_X)$$

into $\mathcal{P}^*$, and $\tau^* \upharpoonright \lambda_X = \sigma \upharpoonright \lambda_X$.

Proof. Immediate from the definitions of $\mathcal{P}^*$ and τ , clause (c) of Lemma 2.4.2, and Łoś' theorem. $\square$

Lemma 2.4.4. $\mathcal{M}(\mathcal{S}_Y, \infty)$ and $\mathcal{P}^*$ agree below λ .

Proof. Clearly $\mathcal{M}(\mathcal{S}_Y, \infty)$, $\mathcal{M}(\mathcal{S}, \infty)$, and $\mathcal{M}(\mathcal{T}, \infty)$ all agree beyond λ . By coherence, $\mathcal{M}(\mathcal{S}_Y, \infty)$ and $\text{ult}(\mathcal{P}_Y, F)$ agree below λ . Clearly $\mathcal{M}(\mathcal{S}_Y, \infty)$ and $\mathcal{P}_Y$ agree below κ . Hence, by clause (d) of Definition 2.1, $\mathcal{M}(\mathcal{S}_Y, \infty)$ and $j(\mathcal{J}_\kappa^{\mathcal{P}_Y})$ agree below λ . By clause (b) of Lemma 2.4.2, $\mathcal{P}_u$ agrees with $\mathcal{J}_\kappa^{\mathcal{P}}$ beyond λ_u for G_b -almost every $u \in B$. It follows by Łoś' theorem that $\mathcal{M}(\mathcal{S}_Y, \infty)$ agrees with $\mathcal{P}^*$ below λ . $\square$

Lemma 2.4.5. *$\mathcal{P}^*$ is conditionally $(\lambda + 1)$ -strong. Moreover, if $\mathcal{S}^*$ is an iteration tree which witnesses that $\mathcal{P}^*$ is conditionally $(\lambda + 1)$ -strong, then $\mathcal{S}^*$ uses the same extenders as $\mathcal{S}_Y$ below λ .*

Proof. By Łoś' theorem and clause (a) of Lemma 2.4.2, $\mathcal{P}^*$ is conditionally $(\lambda + 1)$ -strong in $\text{ult}(N, G)$. By clause (c) of Definition 2.1, $V_{\nu+1} \subset \text{ult}(N, G)$. It follows that $\mathcal{P}^*$ is conditionally $(\lambda + 1)$ -strong in V . In light of Lemma 2.4.4, any iteration tree $\mathcal{S}^*$ witnessing that $\mathcal{P}^*$ is conditionally $(\lambda + 1)$ -strong must use the same extenders as $\mathcal{S}_Y$ below λ . $\square$

Lemma 2.4.6. *$\Phi(\mathcal{S}_Y) \frown \langle \mathcal{P}^*, k, \nu \rangle$ is an iterable phalanx.*

Proof. Put $\mathfrak{A} = \Phi(\mathcal{S}_Y) \frown \langle \mathcal{P}^*, k, \nu \rangle$. That $\mathfrak{A}$ is a phalanx follows from Lemma 2.4.4.

Fix $\mathcal{S}^*$ and an embedding ψ of $\mathcal{P}^*$ into an initial segment of $\mathcal{M}(\mathcal{S}^*, \infty)$ which witness that $\mathcal{P}^*$ is conditionally $(\lambda + 1)$ -strong.

Let us write $\psi(\mathcal{P}^*)$ for the initial segment of $\mathcal{M}(\mathcal{S}^*, \infty)$ into which ψ elementarily embeds $\mathcal{P}^*$. Since $\mathcal{M}(\mathcal{S}^*, 0)$ is $(\lambda + 1)$ -strong, there is

- a successor-length iteration tree $\mathcal{R}$ on W , such that $\text{lh}(E_\eta^{\mathcal{R}}) > \lambda$ whenever $\eta < \text{lh}(\mathcal{R})$, and
- an elementary embedding φ of $\mathcal{M}(\mathcal{S}^*, 0)$ into an initial segment of $\mathcal{M}^{\mathcal{R}}_\infty$ such that $\text{crit}(\varphi) > \lambda$.

We may use φ to copy $\mathcal{S}^*$ to an iteration tree $\varphi\mathcal{S}^*$ on $\mathcal{M}^{\mathcal{R}}_\infty$. Let

$$\varphi_\infty : \mathcal{M}(\mathcal{S}^*, \infty) \longrightarrow \mathcal{M}(\varphi\mathcal{S}^*, \infty)$$

be the final map in the aforementioned copying construction. Then $\text{crit}(\varphi_\infty) > \lambda$ and $\varphi\mathcal{S}^*$ uses the same extenders as $\mathcal{S}_Y$ below λ . Therefore $\varphi_\infty \circ \psi$ is an elementary embedding of $\mathcal{P}^*$ into the initial segment $(\varphi_\infty \circ \psi)(\mathcal{P}^*)$ of $\mathcal{M}(\varphi\mathcal{S}^*, \infty)$ and

$$\text{crit}(\varphi_\infty \circ \psi) > \lambda.$$

Recall that $\mathcal{T}$ was our original iteration tree on W . Put

$$\mathfrak{B} = \Phi(\mathcal{T}) \frown \langle (\varphi_\infty \circ \psi)(\mathcal{P}^*), k, \nu \rangle$$

The agreement described above implies that $\mathfrak{B}$ is a phalanx. It is a phalanx “derived” from $\mathcal{T}$ and $\mathcal{R} \frown \varphi\mathcal{S}^*$, both iteration trees on W . Such W -generated phalanxes are iterable by [St1, 6.9].

But now, the pair of maps $(\pi_Y, (\varphi_\infty \circ \psi))$ can be used in the standard way to reduce the iterability of $\mathfrak{A}$ to that of $\mathfrak{B}$. $\square$

Recall that $\mathcal{V}_X$ is a simple, illbehaved iteration tree on

$$\Phi(\mathcal{S}_X) \frown \langle \text{ult}_k(\mathcal{P}_X, F_X), k, \nu_X \rangle$$

Using the pair of maps (σ, τ^*) we can copy $\mathcal{V}_X$ to an illbehaved iteration tree on $\Phi(\mathcal{S}_Y) \smallfrown \langle \mathcal{P}^*, k, \nu \rangle$. But now we have a contradiction of Lemma 2.4.6. This completes the proof of Theorem 2.3. $\square$

We conclude this section with a version of Theorem 2.3 for mice with sufficiently small projectum. Note that, unlike in Theorem 2.3, we do not assume that there is no inner model with a Woodin cardinal, nor any other smallness condition, in Theorem 2.5.

Theorem 2.5. *Let $\mathcal{M}$ be a sound premouse such that*

$$\rho_\omega(\mathcal{M}) \leq \text{crit}(E)$$

whenever E is an extender from the $\mathcal{M}$ -sequence. Suppose that every countable phalanx which is realizable in an $\mathcal{M}$ -based phalanx is ω_1 -iterable. Let $\mathcal{T}$ be a normal iteration tree on $\mathcal{M}$. Suppose that F is an extender that coheres with $\mathcal{M}_\infty^\mathcal{T}$ such that

$$\text{lh}(E_\eta^\mathcal{T}) < \text{lh}(F)$$

for all $\eta < \text{lh}(\mathcal{T})$, and such that

$$\rho_\omega(\mathcal{M}) < \text{crit}(F).$$

Suppose that $(\mathcal{M}_\infty^\mathcal{T}, F)$ is weakly countably certified. Then F is on the $\mathcal{M}_\infty^\mathcal{T}$ -sequence.

Proof. Put $\kappa = \text{crit}(F)$, $\nu = \nu(F)$, and $\lambda = \text{lh}(F)$. Let θ be a large regular cardinal and $\pi : N \longrightarrow V_\theta$ be the uncollapse of a countable elementary substructure of V_θ . Say $\pi(\overline{\mathcal{T}}) = \mathcal{T}$, $\pi(\overline{F}) = F$, $\pi(\overline{\nu}) = \nu$, and $\pi(\overline{\lambda}) = \lambda$. By the argument in the proof of Theorem 2.3, there is an iteration tree $\mathcal{T}^*$ on $\mathcal{M}$ such that $\mathcal{M}(\mathcal{T}^*, \infty)$ agrees with $\mathcal{M}_\alpha^\mathcal{T}$ below ν whenever $\alpha < \text{lh}(\mathcal{T})$ and a map τ from $\mathcal{M}(\overline{\mathcal{T}}, \infty)$ into $\mathcal{M}(\mathcal{T}^*, \infty)$ such that $\tau \upharpoonright \lambda = \pi \upharpoonright \lambda$. Moreover, if k is the degree of the F -extension of $\Phi(\mathcal{T})$, then τ is a weak k -embedding which is Σ_{k+1} -elementary. This implies that the $\overline{F}$ -extension of $\Phi(\overline{\mathcal{T}})$ is realizable in the $\mathcal{M}$ -based phalanx

$$\Phi(\mathcal{T}) \smallfrown \langle \mathcal{M}(\mathcal{T}^*, \infty), k, \nu \rangle.$$

By the iterability hypothesis on $\mathcal{M}$, there is a successful coiteration $(\mathcal{U}, \mathcal{V})$ of $\Phi(\overline{\mathcal{T}})$ and the $\overline{F}$ -extension of $\Phi(\overline{\mathcal{T}})$. Just as in the proof of [St1, 8.6], our hypothesis on the projectum of $\mathcal{M}$ allows us to conclude that $\overline{F}$ is on the $\mathcal{M}(\overline{\mathcal{T}}, \infty)$ -sequence. (Note that we do not know that $(\mathcal{U}, \mathcal{V}) \in N$ until after arguing that $\overline{F}$ is on the $\mathcal{M}(\overline{\mathcal{T}}, \infty)$ -sequence.) By the elementarity of π , F is on the $\mathcal{M}_\infty^\mathcal{T}$ -sequence. $\square$

3. APPLICATIONS OF MAXIMALITY

Kunen showed that if κ is a weakly compact cardinal and

$$(\kappa^+)^L < \kappa^+,$$

then $0^\#$ exists; see [Jech, p.384]. Our first application of Theorem 2.3 is an extension of Kunen's result to the current generation of core models.

Theorem 3.1. *Suppose that Ω is a measurable cardinal, and that there is no inner model with a Woodin cardinal. Let κ be a weakly compact cardinal $< \Omega$. Then $(\kappa^+)^K = \kappa^+$.*

The result analogous to Theorem 3.1 was proved for measurable cardinals κ in [St1, 8.15], for countably closed, singular cardinals in [MiSchSt], and for all singular cardinals in [MiSch]. From [MiSch], we already know that $(\kappa^+)^K$ has cofinality at least κ whenever κ is a cardinal $\geq \aleph_2$.

Proof of Theorem 3.1. We shall write K^A for $K^{L[A]}$ for any set A . Our first few lemmas will be used in later sections and do not require that κ be a weakly compact cardinal.

Lemma 3.1.1. *Let κ be a cardinal. Suppose that A is a bounded subset of Ω such that $(H_\kappa)^{L[A]} = H_\kappa$. Then K^A and K agree below κ . Moreover, for any K -cardinal $\alpha < \kappa$ and any properly small premouse $\mathcal{P} \in L[A]$:*

$$\mathcal{P} \text{ is } \alpha\text{-strong} \iff (\mathcal{P} \text{ is } \alpha\text{-strong})^{L[A]}.$$

For the definitions, see [St1, 6.4, 6.12]. To prove Lemma 3.1.1, one must note that Ω is an $L[A]$ -indiscernible, and that all the theorems of [St1] hold for K constructed inside $L[A]$. Then Lemma 3.1.1 is immediate from the following minor strengthening of [St1, 6.14].

Lemma 3.1.2. *Suppose that α is a cardinal of K . Let $\mathcal{P}$ be a properly small premouse such that $\mathcal{J}_\alpha^\mathcal{P} = \mathcal{J}_\alpha^K$, but $\mathcal{P}$ is not α -strong. Then there exists a properly small premouse $\mathcal{Q}$ which is β -strong for every $\beta < \alpha$, such that*

$$\text{card}(|\mathcal{Q}|) = \text{card}(\alpha),$$

and there exists a countable, simple, illbehaved iteration tree $\mathcal{T}$ on $((\mathcal{Q}, \mathcal{P}), \alpha)$.

The only difference between Lemma 3.1.2 and [St1, 6.14] is that we do not assume that $\text{card}(|\mathcal{P}|) = \text{card}(\alpha)$. But the proof of [St1, 6.14] gives this with no additional work. Since Lemma 3.1.2 holds, not only in V , but also in any $L[A]$, Lemma 3.1.1 follows.

Lemma 3.1.3. *Suppose that κ is a cardinal of uncountable cofinality, and let $\mathcal{P}$ be a properly small premouse with $\mathcal{J}_\kappa^\mathcal{P} = \mathcal{J}_\kappa^K$. Then:*

$$\mathcal{P} \text{ is } \kappa\text{-strong} \iff \forall K\text{-cardinal } \alpha < \kappa \text{ } (\mathcal{P} \text{ is } \alpha\text{-strong}).$$

Proof. The $\implies$ direction is obvious from the definition; see [St1, 6.4]. So suppose that $\mathcal{P}$ is not κ -strong. By Lemma 3.1.2, there is an κ -strong, properly small premouse $\mathcal{Q}$ of cardinality κ such that $\mathcal{J}_\kappa^\mathcal{Q} = \mathcal{J}_\kappa^K$, and a countable, illbehaved iteration tree $\mathcal{T}$ on $((\mathcal{Q}, \mathcal{P}), \kappa)$. Since $\mathcal{T}$ is countable and κ has uncountable cofinality, there is an $\alpha < \kappa$ such that $\mathcal{T}$ can be construed as an iteration tree on the phalanx $((\mathcal{Q}, \mathcal{P}), \alpha)$. But then $\mathcal{P}$ is not α -strong by [St1, 6.11]. $\square$

Also, Lemma 3.1.3 holds in $L[A]$ for any set $A \in V_\Omega$. Directly from [St1, 6.4], we see that both V and $L[A]$ satisfy the sentence: “Every properly small, κ -strong premouse $\mathcal{P}$ with $\mathcal{J}_\kappa^\mathcal{P} = \mathcal{J}_\kappa^K$ and $\rho_\omega(\mathcal{P}) = \kappa$ is a level of K , and for cofinally many $\gamma < (\kappa^+)^K$, $\mathcal{J}_\gamma^K$ is a properly small, κ -strong premouse which projects to κ .” But then, it follows that:

Lemma 3.1.4. *Let κ be a cardinal of uncountable cofinality. Suppose that A is a bounded subset of Ω such that $(H_\kappa)^{L[A]} = H_\kappa$. Then K^A and K agree below $(\kappa^+)^{K^A}$. Moreover, for any properly small premouse $\mathcal{P} \in L[A]$:*

$$\mathcal{P} \text{ is } \kappa\text{-strong} \iff (\mathcal{P} \text{ is } \kappa\text{-strong})^{L[A]}.$$

The hypothesis of uncountable cofinality is not needed in Lemma 3.1.4, as we shall show in Lemma 4.2.2.

Now fix a weakly compact cardinal $\kappa < \Omega$ and put $\lambda = (\kappa^+)^K$. Assume, for contradiction, that $\lambda < \kappa^+$.

Let N be a transitive, power admissible set such that ${}^\omega N \subset N$,

$$V_\kappa \cup \mathcal{J}_{\lambda+1}^K \subset N,$$

and $\text{card}(N) = \kappa$. Let $A \subseteq \kappa$ such that $N \in L[A]$.

It follows immediately from Lemma 3.1.4 that $\lambda = (\kappa^+)^{K^A}$, and that K^A and K agree below λ .

Since $A^\#$ exists, $(\kappa^+)^{L[A]} < \kappa^+$, so $\text{card}(\mathcal{P}(\kappa) \cap L[A]) = \kappa$. Because κ is weakly compact, there is a non-principal, κ -complete $L[A]$ -ultrafilter U over κ ; see [Jech, p.384]. Then $\text{ult}(L[A], U)$ is well-founded. Let

$$j : L[A] \longrightarrow \text{ult}(L[A], U) = L[j(A)]$$

be the ultrapower map. Then $\text{crit}(j) = \kappa$ and $A = j(A) \cap \kappa \in L[j(A)]$, so $L[A] \subseteq L[j(A)]$. We remark that $\mathcal{P}(\kappa) \cap L[j(A)] \not\subseteq L[A]$ is possible;

see [Jech, p.394]. It follows from Lemma 3.1.4 that $\lambda = (\kappa^+)^{K^{j(A)}}$, and that $K^{j(A)}$ and K agree past λ . Thus $\mathcal{P}(\kappa) \cap K^{j(A)} = \mathcal{P}(\kappa) \cap K^A$.

Let E_j be the superstrong extender derived from j . So, for every $a \in [j(\kappa)]^{<\omega}$,

$$(E_j)_a = \{x \subseteq [\kappa]^{|a|} \mid x \in L[A] \wedge a \in j(x)\}.$$

We claim that for every sequence $\langle x_\alpha \mid \alpha < \kappa \rangle$ in $L[A]$,

$$E_j \cap ([j(\kappa)]^{<\omega} \times \{x_\alpha \mid \alpha < \kappa\}) \in L[j(A)].$$

The argument is due to Kunen, as in [MiSt, 1.1]: just note that for any $a \in [j(\kappa)]^{<\omega}$,

$$E_a = \{x_\alpha \subseteq [\kappa]^{|a|} \mid \alpha < \kappa \wedge a \in j(x_\alpha)\}$$

and

$$\langle j(x_\alpha) \mid \alpha < \kappa \rangle \in L[j(A)].$$

Set

$$F = E_j \cap ([j(\kappa)]^{<\omega} \times K^A)$$

and

$$G = E_j \cap ([j(\kappa)]^{<\omega} \times N).$$

Then $(K^{j(A)}, F)$ and (N, G) are elements of $L[j(A)]$. Moreover, $L[j(A)]$ satisfies the sentence, “ $F \restriction \xi$ coheres with K and (N, G) is an $\mathcal{A}$ -certificate for $(K, F \restriction \xi)$ ” where

$$\mathcal{A} = \bigcup_{n < \omega} \mathcal{P}([\kappa]^n)^K$$

and unbounded many $\xi < j(\kappa)$. (For clause (c) of Definition 2.1, simply note that $L[A] \models “V_\kappa \subset N”$, so

$$L[j(A)] \models “V_{j(\kappa)} \subset j(N)”$$

and $j(N)$ and $\text{ult}(N, G)$ have the same sets of rank $< j(\kappa)$.) That Ω is an $L[j(\kappa)]$ -indiscernible is enough to see that the conclusion of Theorem 2.3 holds in $L[j(\kappa)]$. Therefore every initial segment of F is on the $K^{j(A)}$ sequence. But then κ is a Shelah cardinal in $K^{j(A)}$, contradicting that there is no inner model with a Woodin cardinal. $\square$

By definition, K is the transitive collapse of an elementary substructure of K^c ; see [St1, §5]. Our next application of Theorem 2.3 shows that there is an iteration tree on K with last model K^c . For the statement of Theorem 3.2, recall that K^c is defined as the limit of the maximal 1-small construction $\langle \mathcal{N}_\xi \mid \xi < \Omega \rangle$; see [St1, §1].

Theorem 3.2. *Assume that Ω is a measurable cardinal and that there is no inner model with a Woodin cardinal. Let $\alpha \leq \Omega$ and $k < \omega$. Suppose that $(\mathcal{U}, \mathcal{V})$ is the coiteration of*

$$(K, \mathfrak{C}_k(\mathcal{N}_\xi)).$$

Then $\mathcal{V}$ is trivial. In particular, since K is universal, K^c is an iterate of K .

Proof. The proof is by induction. Fix $\xi \leq \Omega$ and $k < \omega$. Assume that $\mathfrak{C}_\ell(\mathcal{N}_\zeta)$ does not move in its coiteration with K whenever

$$(\zeta, \ell) <_{lex} (\xi, k).$$

For contradiction, assume that $\mathfrak{C}_k(\mathcal{N}_\xi)$ moves in its coiteration with K .

Let $(\mathcal{U}, \mathcal{V})$ be the coiteration of

$$(K, \mathfrak{C}_k(\mathcal{N}_\xi))$$

and $\theta < \text{lh}(\mathcal{V})$ be least such that $E_\theta^\mathcal{V} \neq \emptyset$. Put $\mathcal{T} = \mathcal{U} \restriction (\theta + 1)$ and $F = E_\theta^\mathcal{V}$. Then F coheres with $\mathcal{M}_\theta^\mathcal{T}$ but F is not on the $\mathcal{M}_\theta^\mathcal{T}$ -sequence.

First suppose that $k = 0$ and that F is the last extender of $\mathcal{N}_\xi$. Then F is countably certified and $\text{crit}(F)$ is an inaccessible cardinal, by the definition of $\mathcal{N}_\xi$. Hence, by Theorem 3.2, F is on the $\mathcal{M}_\theta^\mathcal{T}$ -sequence, a contradiction.

Next, suppose that $k = 0$ but that F is not the last extender of $\mathcal{N}_\xi$. Then there is some $\zeta < \xi$ and $\ell < \omega$ such that F is on the $\mathfrak{C}_\ell(\mathcal{N}_\zeta)$ -sequence. From the fact that $\mathfrak{C}_\ell(\mathcal{N}_\zeta)$ and $\mathfrak{C}_k(\mathcal{N}_\xi)$ agree below $\text{lh}(F) + 1$, it follows that $\mathfrak{C}_\ell(\mathcal{N}_\zeta)$ moves in its coiteration with K . This contradicts our induction hypothesis.

So $k > 0$. By the induction hypothesis,

$$\mathfrak{C}_{k-1}(\mathcal{N}_\xi) \neq \mathfrak{C}_k(\mathcal{N}_\xi).$$

Thus, $\mathfrak{C}_{k-1}(\mathcal{N}_\xi)$ is $(k-1)$ -sound but not k -sound. By the induction hypothesis, there is an iteration tree $\mathcal{S}$ on K such that the last model of $\mathcal{S}$ is $\mathfrak{C}_{k-1}(\mathcal{N}_\xi)$. But then $\mathfrak{C}_k(\mathcal{N}_\xi)$ is a model along the main branch of $\mathcal{S}$. Namely,

$$\mathfrak{C}_k(\mathcal{N}_\xi) = (\mathcal{M}_{\gamma+1}^*)^\mathcal{S}$$

where $\gamma + 1$ is the last drop to an ultrapower of degree at most $k - 1$ along the main branch of $\mathcal{S}$. But then $\mathcal{S} \restriction (\gamma + 1)$ witnesses that the theorem holds for (ξ, k) , a contradiction. $\square$

Recall that $\mathcal{N}_\kappa$ is a premouse of height κ whenever κ is a cardinal, but that $\mathcal{N}_\kappa$ need not equal $\mathcal{J}_\kappa^{K^c}$. Theorem 3.3 appears to be the first step in showing that $\mathcal{N}_\kappa = \mathcal{J}_\kappa^{K^c}$ when κ is a Mahlo cardinal, but we do not know if this is true.

Theorem 3.3. *Suppose that there is no inner model with a Woodin cardinal and that Ω is a measurable cardinal. Let $\kappa < \Omega$ be a Mahlo cardinal. Then there is an iteration tree $\mathcal{T}$ on $\mathcal{J}_\kappa^K$ such that $\mathcal{M}_\infty^\mathcal{T} = \mathcal{N}_\kappa$. In particular,*

$$(\alpha^+)^{\mathcal{N}_\kappa} = \alpha^+$$

for all singular cardinals $\alpha < \kappa$, so $\mathcal{N}_\kappa$ is universal for mice of height $\leq \kappa$.

Proof. By Theorem 3.2, there is an iteration tree $\mathcal{T}$ on K with $\mathcal{M}_\infty^\mathcal{T} = \mathcal{N}_\kappa$. Certainly $\text{lh}(\mathcal{T}) \leq \kappa + 1$. Let us assume that $\text{lh}(\mathcal{T}) = \kappa + 1$, the other case being clear. It is enough if we show that $\mathcal{T}$ does not drop and that $i_{0,\kappa}^\mathcal{T}(\alpha) < \kappa$ for all $\alpha < \kappa$.

Suppose otherwise. Form an internally approachable elementary chain

$$\langle X_\alpha \mid \alpha < \kappa \rangle$$

of submodels of $\langle V_{\kappa+1}, \in, \mathcal{T} \rangle$ such that $\text{card}(X_\alpha) = \alpha$ and ${}^\omega X_\alpha \in X_{\alpha+1}$ for all $\alpha < \kappa$. Let

$$\pi_\alpha : P_\alpha \longrightarrow V_{\kappa+1}$$

be the uncollapse of X_α and G_α be the length κ extender derived from π_α whenever $\alpha < \kappa$. There is a stationary set of inaccessible cardinals $\alpha < \kappa$ such that if $(\beta+1)T_\kappa$ and $\alpha = \text{pred}^\mathcal{T}(\beta+1)$, then $\alpha = \text{crit}(\pi_\alpha)$, $\pi_\alpha(\alpha) = \kappa$, X_α is countably closed, $\mathcal{M}_\alpha^\mathcal{T} \cap \mathcal{P}(\alpha) \in P_\alpha$, and (P_α, G_α) is an $\mathcal{A}$ -certificate for $(\mathcal{M}_\beta^\mathcal{T}, E_\beta^\mathcal{T})$ where

$$\mathcal{A} = \bigcup_{n < \omega} \mathcal{P}([\alpha]^n) \cap |\mathcal{M}_\beta^\mathcal{T}|.$$

The proof is similar to the standard argument that coiterations of mice terminate. Fix any such $\alpha < \kappa$. Clearly $E_\beta^\mathcal{T}$ coheres with $\mathcal{N}_\kappa$ and $(\mathcal{N}_\kappa, E_\beta^\mathcal{T})$ is countably certified. Note that $\text{lh}(E_\beta^\mathcal{T})$ is a cardinal in $\mathcal{N}_\kappa$. Let $\gamma < \kappa$ be the supremum of the $\eta < \kappa$ such that $\rho_\omega(\mathcal{N}_\eta) < \text{lh}(E_\beta^\mathcal{T})$. It is not hard to see that $\mathcal{N}_\gamma$ is the passive initial segment of $\mathcal{N}_\kappa$ of height $\text{lh}(E_\beta^\mathcal{T})$, $\mathcal{N}_\gamma \cap \langle E_\beta^\mathcal{T} \rangle$ is a premouse, and that $(\mathcal{N}_\gamma, E_\beta^\mathcal{T})$ is certified as above. Therefore

$$\rho_\omega(\mathcal{N}_{\gamma+1}) \leq \nu(E_\beta^\mathcal{T}) < \text{lh}(E_\beta^\mathcal{T}),$$

a contradiction. □

Theorem 3.4 (with Mitchell). *Suppose that there is no inner model with a Woodin cardinal and that Ω is a measurable cardinal. Let κ be a cardinal such that $\aleph_2 \leq \kappa < \Omega$. Then $\mathcal{J}_\kappa^K$ is universal for premice of height κ .*

Theorem 3.4 reduces quickly to the case in which κ is a successor cardinal. The authors proved Theorem 3.4 under the assumption that κ is a countably closed, successor cardinal. Then, Mitchell saw how to eliminate the countable closure, by adapting the methods of [MiSch].

Sketch of Theorem 3.4. Suppose that $\mathcal{M}$ is a counterexample to the theorem and that $(\mathcal{S}, \mathcal{T})$ is the coiteration of $(\mathcal{J}_\kappa^K, \mathcal{M})$. Then, there is club set $C \subseteq [0, \kappa]_\mathcal{S}$ such that

$$\begin{aligned} \mathcal{D}^\mathcal{T} \cap [\alpha, \kappa]_T &= \emptyset, \\ \text{crit}(i_{\alpha, \kappa}^\mathcal{T}) &= \alpha, \end{aligned}$$

and

$$i_{\alpha, \kappa}^\mathcal{T}(\alpha) = \kappa$$

whenever $\alpha \in C$.

Obtain (P_α, G_α) from an internally approachable elementary chain of length κ as in the proof of Theorem 3.3. Here, $\text{card}(P_\alpha) = \mu$ and it is possible that no P_α is countably closed.

Assuming that $\mu^\omega < \kappa$, it is possible to arrange countable closure. Then, as in the proof of Theorem 3.3, for a stationary subset of C , (P_α, G_α) is a weak $\mathcal{A}$ -certificate for $(\mathcal{M}_\alpha^\mathcal{T}, E_\alpha^\mathcal{T})$ where

$$\mathcal{A} = \bigcup_{n < \omega} \mathcal{P}([\alpha]^n) \cap |\mathcal{M}_\beta^\mathcal{T}|.$$

It follows that $(\mathcal{M}_\alpha^\mathcal{T}, E_\alpha^\mathcal{T})$ is weakly countably certified. (Note that (P_α, G_α) cannot be an $\mathcal{A}$ -certificate since $V_\alpha \not\subseteq P_\alpha$.) By Theorem 2.3, $E_\alpha^\mathcal{T}$ is on the $\mathcal{M}_\alpha^\mathcal{S}$ -sequence, an obvious contradiction.

Now suppose that $\mu^\omega \geq \kappa$. The only difference is that, with the same definitions as above, we cannot arrange that P_α is countably closed. But notice that in the proof of Theorem 2.3, countable closure was really only used to see that the $E_\alpha^\mathcal{T}$ -extension of $\Phi(\mathcal{S})$ is iterable (so that we could apply [St1, 8.6]). The method of [MiSch] adapts to show that for a stationary set of $\alpha < \kappa$, the $E_\alpha^\mathcal{T}$ -extension of $\Phi(\mathcal{S})$ is iterable. $\square$

4. SQUARE PRINCIPLES

Recall from [Sch1, 5.1] or [Sch2] the following hierarchy of principles intermediate to Jensen's well-known principles $\square_\kappa$ and $\square_\kappa^*$ from [J1]. If $1 < \lambda \leq \kappa^+$, then $\square_\kappa^{<\lambda}$ is the principle asserting the existence of a sequence $\langle \mathcal{F}_\nu \mid \nu < \kappa^+ \rangle$ such that for every limit ordinal $\nu \in (\kappa, \kappa^+)$,

- (1) $1 \leq \text{card}(\mathcal{F}_\nu) < \lambda$, and
- (2) for all $C \in \mathcal{F}_\nu$,
 - (a) C is a closed, unbounded subset of ν ,

- (b) C has order type $\leq \kappa$, and
- (c) $C \cap \mu \in \mathcal{F}_\mu$ whenever $\mu < \nu$ is a limit point of C .

We write $\square_\kappa^\lambda$ for $\square_\kappa^{<\lambda^+}$; so $\square_\kappa \equiv \square_\kappa^1$ and $\square_\kappa^* \equiv \square_\kappa^\kappa$. Using these forms of weak square, some lower bounds on the large cardinal consistency strength of the Proper Forcing Axiom and other principles were obtained in [Sch1, Sch2, SchSt].

Theorem 3.1 and its proof will be used to prove Theorem 4.1 below.

Theorem 4.1. *Suppose that there is no inner model with a Woodin cardinal. Then $\square_\kappa^{<\omega}$ holds whenever κ is a weakly compact cardinal.*

If there is a measurable cardinal Ω , but no inner model with a Woodin cardinal, then $\square_\kappa^{<\omega}$ holds for every weakly compact cardinal $\kappa < \Omega$; this is immediate from Theorem 3.1 and [Sch2]. Theorem 4.1 says that the measurable cardinal is not necessary.

Our proof of Theorem 4.1 can be modified to show that if $\square_\kappa^{<\omega}$ fails, then every $x \subseteq \kappa$ is in a proper class inner model $M_1(x)$ with a Woodin cardinal, and that $M_1(x)^\#$ exists. It follows from unpublished work of Woodin that $\neg \square_\kappa^{<\omega}$ implies Σ_2^1 -determinacy; see [N] for related results. Using our proof of Theorem 4.1, Woodin has since shown that $\neg \square_\kappa^{<\omega}$ implies $L(\mathbb{R})$ -determinacy. Woodin's proof is an induction on the levels of $L(\mathbb{R})$, with Theorem 4.1 as the base case, and he makes the general case look like the base case. Woodin had used this method before, in unpublished work.

Proof of Theorem 4.1. Suppose that κ is a weakly compact cardinal and that there is no inner model with a Woodin cardinal. Assume for contradiction that $\square_\kappa^{<\omega}$ fails. Then $A^\#$ exists whenever A is a bounded subset of κ . This follows from Jensen's theorem that $\square_\kappa$ holds in $L[A]$ (cf. [J1]) and Kunen's theorem on successors of weakly compact cardinals relativized to $L[A]$ (cf. [Jech, p.384]). Since κ is weakly compact, $A^\#$ exists for all $A \subseteq \kappa$. Let $\Omega = \kappa^+$. Then Ω is an A -indiscernible for every $A \subseteq \kappa$. So, for every $A \subseteq \kappa$, the theory of K up to Ω of [St1] applies in $L[A]$.

Let $\langle A_\alpha \mid \alpha < \Omega \rangle$ be a sequence of subsets of κ and $\langle \lambda_\alpha \mid \alpha < \Omega \rangle$ be a sequence of ordinals with $A_\alpha \in L[A_\beta]$,

$$(\kappa^+)^{L[A_\alpha]} = \lambda_\alpha \geq \alpha ,$$

and $\lambda_\alpha < \lambda_\beta$ whenever $\alpha < \beta < \Omega$. Also, assume that $V_\kappa^{L[A_\alpha]} = V_\kappa$.

Note that κ is weakly compact in $L[A_\alpha]$ whenever $\alpha < \Omega$. Therefore,

$$\lambda_\alpha = (\kappa^+)^{K^{A_\alpha}}$$

whenever $\alpha < \Omega$, by a version of Theorem 3.1 applied inside $L[A_\alpha]$. By Lemma 3.1.4, K^{A_α} and K^{A_β} agree below λ_α whenever $\alpha < \beta < \Omega$. Let

$$\vec{E} = \bigcup_{\alpha < \Omega} \left(\dot{E}^{K^{A_\alpha}} \restriction \lambda_\alpha \right)$$

and $\mathcal{M} = \mathcal{J}_\Omega^{\vec{E}}$. Clearly,

$$(\kappa^+)^{\mathcal{M}} = \Omega.$$

By [Sch2], $\square_\kappa^{<\omega}$ holds in K^{A_α} for all $\alpha < \Omega$. The proof in [Sch2] is “local” and gives more. Namely, for each $\alpha < \Omega$, there a sequence $\mathcal{F}^\alpha = \langle \mathcal{F}_\nu^\alpha \mid \nu < \lambda_\alpha \rangle$ that witnesses $\square_\kappa^{<\omega}$ in K^{A_α} , such that $\mathcal{F}^\alpha = \mathcal{F}^\beta \restriction \lambda_\alpha$ whenever $\alpha < \beta < \Omega$. So $\mathcal{F} = \bigcup_{\alpha < \Omega} \mathcal{F}^\alpha$ witnesses $\square_\kappa^{<\omega}$ in $\mathcal{M}$, and therefore in V . This is a contradiction. $\square$

If there is a measurable cardinal Ω , but no inner model with a Woodin cardinal, then $\square_\kappa^{<\omega}$ holds for every singular cardinal $\kappa < \Omega$. This is immediate from [MiSch] and [Sch2]. The next theorem says that the measurable cardinal is not necessary, at least when κ is a strong limit cardinal.

Theorem 4.2. *Suppose that there is no inner model with a Woodin cardinal. Then $\square_\kappa^{<\omega}$ holds whenever κ is a singular, strong limit cardinal.*

Jensen proved that if $0^\sharp$ does not exist, then $\square_\kappa$ holds whenever κ is a singular cardinal. (Cf. [J2] and [J3].) It is not known if Theorem 4.2 holds for all singular cardinals, nor if its conclusion can be strengthened to $\square_\kappa$ holding. Woodin’s inductive proof of determinacy mentioned before adapts to show that $\neg \square_\kappa^{<\omega}$ implies Inductive Determinacy, hence Projective Determinacy. (Using the lower-part closure Lp instead of L in the proof, one can see directly that if $\square_\kappa^{<\omega}$ fails and κ is a singular, strong limit cardinal, then, over any bounded subset of κ , there is a premouse $\mathcal{M}$ with ω -many Woodin cardinals cofinal in $\text{OR}^{\mathcal{M}}$.) It is not known if $\neg \square_\kappa^{<\omega}$ implies $L(\mathbb{R})$ -determinacy.

Proof of Theorem 4.2. Suppose that κ is a singular, strong limit cardinal and that there is no inner model with a Woodin cardinal. Assume for contradiction that $\square_\kappa^{<\omega}$ fails. Jensen showed that $\square_\kappa$ holds in $L[A]$ whenever $A \subseteq \kappa$ (cf. [J1]). Therefore $(\kappa^+)^{L[A]} < \kappa^+$ whenever $A \subseteq \kappa$.

Lemma 4.2.1. *$A^\#$ exists whenever $A \subseteq \kappa$.*

Sketch. It is clear that Jensen’s covering lemma (cf. [DeJ]) can be adapted to show that $A^\#$ exists whenever A is a bounded subset of κ . Suppose that A is an unbounded subset of κ and that $\lambda = (\kappa^+)^{L[A]} < \kappa$.

κ^+ . We shall get $A^\#$ by piecing together lift-ups of sharps for bounded subsets of κ .

Let $\pi : N \longrightarrow V_\theta$ for some large θ with N transitive, countably closed, $\text{card}(N) < \kappa$, and π cofinal in λ . Say $\pi(\bar{A}) = A$, $\pi(\bar{\kappa}) = \kappa$, and $\pi(\bar{\lambda}) = \lambda$. Let E be the length κ extender derived from π . If $\bar{\lambda}$ is not a cardinal of $L[\bar{A}]$, then as usual we let $\mathcal{M}$ be the first level of $L[\bar{A}]$ which sees this, note that

$$\mathcal{M} = \mathcal{H}_{n+1}^{\mathcal{M}}(\bar{\kappa} \cup p_{n+1}^{\mathcal{M}})$$

for an appropriate $n < \omega$ (fine structure works above $\bar{\kappa}$), set $\mathcal{M}^* = \text{ult}_n(\mathcal{M}, E)$, and get that $\mathcal{M}^*$ is a level of $L[A]$ collapsing λ .

So $\bar{\lambda}$ is a cardinal of $L[\bar{A}]$, and thus E measures all subsets of $\bar{\kappa}$ in $L[\bar{A}]$. Let $\pi^* : L[\bar{A}] \longrightarrow \text{ult}(L[\bar{A}], E) = L[A]$ be the lift-up of π . Let D be the class of strong limit cards of cofinality κ^+ . Note the tuples from D are $\bar{A}$ -indiscernible with respect to parameters $< \bar{\kappa}$. Also, $\pi^*(\alpha) = \alpha$ for $\alpha \in D$. By the elementarity of π^* , the tuples from D are A -indiscernible with respect to parameters from κ . Since D is stationary, we are done. $\square$

Let $\vec{\kappa} = \langle \kappa_i \mid i < \text{cf}(\kappa) \rangle$ be an increasing sequence of cardinals cofinal in κ . Put $\Omega = \kappa^+$. Choose $\langle A_\alpha \mid \alpha < \Omega \rangle$ and $\langle \lambda_\alpha \mid \alpha < \Omega \rangle$ as in the proof of Theorem 4.1, with the additional stipulation that $\vec{\kappa} \in L[A_0]$. Then κ is a singular, strong limit cardinal in $L[A_\alpha]$, so

$$\lambda_\alpha = (\kappa^+)^{K^{A_\alpha}}$$

for all $\alpha < \Omega$, by a version of [MiSchSt] applied inside $L[A_\alpha]$. If κ has uncountable cofinality, then we are done as in the proof of Theorem 4.1. So assume that $\text{cf}(\kappa) = \omega$. Then Lemma 4.2.2 below is clearly enough.

Lemma 4.2.2. *If $\alpha < \beta < \Omega$, then K^{A_α} and K^{A_β} agree below λ_α .*

Proof. Suppose, for contradiction, that the lemma is false. Throughout the proof, we work in $L[A_\beta]$. Put $A = A_\alpha$.

Since K^A and K agree below κ , there is a properly small level $\mathcal{P}$ of K^A such that $\mathcal{P}$ is not κ -strong.

Let $\mathcal{Q}$ be a properly small premouse of cardinality κ such that $((\mathcal{Q}, \mathcal{P}), \kappa)$ is not iterable, and let $\mathcal{T}$ be an illbehaved iteration tree on $((\mathcal{Q}, \mathcal{P}), \kappa)$. By taking elementary substructures, we find a sequence of properly small premice

$$\vec{\mathcal{Q}} = \langle \mathcal{Q}_i \mid i < i < \omega \rangle$$

such that $\mathcal{Q}_i$ is κ_i -strong and $\text{card}(|\mathcal{Q}_i|) = \kappa_i$ whenever $i < \omega$, but the phalanx $((\vec{\mathcal{Q}}, \mathcal{P}), \vec{\kappa})$ is not iterable. Each $\mathcal{Q}_i$ elementarily embeds into $\mathcal{Q}$ with critical point $> \kappa_i$. Moreover, $\mathcal{T}$ can be construed as an

iteration tree on $((\vec{\mathcal{Q}}, \mathcal{P}), \vec{\kappa})$, and thus construed, $\mathcal{T}$ is illbehaved. Let M be the transitive collapse of a countable elementary substructure of V_Ω and let $\pi : M \rightarrow V_\Omega$ be the uncollapse. Assume that

$$\pi \left(((\vec{\mathcal{Q}}^M, \mathcal{P}^M), \vec{\kappa}^M) \right) = ((\vec{\mathcal{Q}}, \mathcal{P}), \vec{\kappa})$$

and $\pi(\mathcal{T}^M) = \mathcal{T}$. By elementarity and absoluteness, $\mathcal{T}^M$ is an illbehaved iteration tree on $((\vec{\mathcal{Q}}^M, \mathcal{P}^M), \vec{\kappa}^M)$.

Now $\mathcal{Q}_i \in L[A]$ for $i < \omega$ since $V_\kappa \subset L[A]$. But there is no reason to believe that $\vec{\mathcal{Q}} \in L[A]$. By [St1, 6.14], $\mathcal{Q}_i$ is κ_i -strong in $L[A]$ whenever $i < \omega$. Because $L[A]$ contains all the reals, $M \cup \{M\} \subset L[A]$.

There is a tree in $L[A]$ that simultaneously searches for:

- a sequence $\vec{\mathcal{R}}$ such that $((\vec{\mathcal{R}}, \mathcal{P}), \vec{\kappa})$ is a phalanx of proper premice and $\mathcal{R}_i$ is κ_i -strong in $L[A]$ whenever $i < \omega$, and
- a sequence $\vec{\sigma}$ of length $\omega + 1$ such that σ_ω is an elementary embedding of $\mathcal{P}^M$ into $\mathcal{P}$ with $\sigma_\omega(\kappa^M) = \kappa$, and σ_i is an elementary embedding of $\mathcal{Q}_i^M$ into $\mathcal{R}_i$ with $\sigma_i(\kappa_i^M) = \kappa_i$ whenever $i < \omega$, such that

$$\sigma_i \restriction (\kappa_i^M + 1) = \sigma_j \restriction (\kappa_i^M + 1)$$

whenever $i < j \leq \omega$.

There is an infinite branch through this tree determined by $\vec{\mathcal{R}} = \vec{\mathcal{Q}}$ and

$$\vec{\sigma} = \langle \pi \restriction \mathcal{Q}_i^M \mid i < \omega \rangle \frown \langle \pi \restriction \mathcal{P}^M \rangle.$$

By the absoluteness, there is a branch through this tree in $L[A]$. So let $\vec{\mathcal{R}}$ and $\vec{\sigma}$ be determined by such a branch.

The iteration tree on $((\vec{\mathcal{R}}, \mathcal{P}), \vec{\kappa})$ obtained by copying $\mathcal{T}^M$ using $\vec{\sigma}$ we denote by $\vec{\sigma}''\mathcal{T}^M$. Now $((\vec{\mathcal{R}}, \mathcal{P}), \vec{\kappa})$ is iterable in $L[A]$ since $\mathcal{P}$ and the $\mathcal{R}_i$ are appropriately strong in $L[A]$. So $\vec{\sigma}''\mathcal{T}$ is well-behaved. Therefore $\mathcal{T}^M$ is well-behaved, a contradiction. $\square$

$\square$

5. AN ERDÖS PROPERTY

It has been observed that the existence of sharps for all sets (rather than a measurable cardinal Ω) suffices to develop the theory in [St1], and consequently, to prove a versions of the results in this paper. Theorem 5.1 below, an improvement to Theorem 3.1, illustrates how an Erdős partition property on Ω consistent with $V = L$ suffices in many core model applications.

Let $\tilde{K}$ be the premouse determined by the inductive definition of the core model in [St1, §6]. By [St1], if Ω is a measurable cardinal, then

$K = \tilde{K}$ up to Ω , but it is not known if one can show in ZFC that $\tilde{K}$ is a proper class.

Theorem 5.1. *Suppose that there is no inner model with a Woodin cardinal. and that $\kappa < \Omega$ are cardinals satisfying the partition relation $\Omega \longrightarrow (\omega)_\kappa^{<\omega}$. Suppose that κ is a weakly compact cardinal. Then $\tilde{K}$ is a premouse of height $> \kappa^+$ and $(\kappa^+)^\tilde{K} = \kappa^+$.*

Of course, Theorem 3.1 is just a special case of Theorem 5.1. The same method of proof can be used to show that the hypothesis

$$\Omega \longrightarrow (\omega)_\kappa^{<\omega}$$

for all $\kappa < \Omega$, is more than enough for all applications in [MiSch], [MiSchSt], [Sch1], [Sch2], [St1], and [St3]. In fact, the proof uses only a model theoretic consequence of the partition relation.

Proof of Theorem 5.1. Suppose that $\kappa < \Omega$ are cardinals such that κ is a weakly compact and

$$\Omega \longrightarrow (\omega)_\kappa^{<\omega}$$

Assume that there is no inner model with a Woodin cardinal.

We shall adapt an argument of Reinhardt and Silver; see [Jech, pp.394–5]. Let $\mathfrak{A}$ be the structure $\langle V_\Omega, \in, \kappa \rangle$ expanded to include predicates for Skolem functions. Let I be an infinite set of indiscernibles for $\mathfrak{A}$ such that $I \subset (\kappa, \Omega)$. The existence of I follows from our assumption that $\Omega \longrightarrow (\omega)_\kappa^{<\omega}$ in the standard way. Let

$$\pi : \mathfrak{B} \longrightarrow \mathfrak{A}$$

be the inverse transitive collapse of the closure of I under the Skolem functions for $\mathfrak{A}$. Fix some $\alpha \in I$ and put $\bar{\alpha} = \pi^{-1}(\alpha)$ and $\bar{\kappa} = \pi^{-1}(\kappa)$. Let $j : \mathfrak{B} \longrightarrow \mathfrak{B}$ be an elementary embedding obtained by “shifting” $\bar{\alpha}$, so that $\text{crit}(j) \leq \bar{\alpha}$. With a judicious choice of I , we can arrange that $\text{crit}(j) > \bar{\kappa}$.

Now, using j just as the embedding from a normal measure was used in [St1], one can show that the inductive definition of the core model, as carried out in $\mathfrak{B}$, succeeds up to $\text{crit}(j)$. The main point is that the superstrong extender from j is weakly amenable to $\mathfrak{B}$. Since π is an elementary embedding, $\tilde{K}$ has height $\geq j(\text{crit}(j)) > \kappa^+$. Arguing as in Theorem 3.1, we see that

$$\mathfrak{B} \models (\bar{\kappa}^+)^K = \bar{\kappa}^+ .$$

By elementarity again, we conclude that $(\kappa^+)^\tilde{K} = \kappa^+$. □

6. THE CORE MODEL IN $L[\vec{E}]$

This section is still in preparation.

REFERENCES

- [DeJ] K.I. Devlin and R.B. Jensen, *Marginalia to a theorem of Silver*, in *Logic Conference, Kiel 1974*, Lecture Notes in Mathematics 499, Springer-Verlag (1975) 115–142.
- [Do1] A.J. Dodd, *Strong Cardinals*, circulated notes (1981).
- [Do2] A.J. Dodd, *The core model*, London Mathematical Society Lecture Note Series 61, Cambridge University Press (1982).
- [DoJ] A.J. Dodd and R.B. Jensen, *The core model*, *Annals of Mathematical Logic* 20 (1981) 43–75.
- [Jech] T. Jech, *Set theory*, Academic Press (1978).
- [J1] R.B. Jensen, *The fine structure of the constructible hierarchy*, *Annals of Mathematical Logic* 4 (1972) 229–308.
- [J2] R.B. Jensen, *Non overlapping extenders*, circulated notes.
- [J3] R.B. Jensen, *Some remarks on $\square$ below $0^\sharp$* , circulated notes.
- [Mi] W.J. Mitchell, *The core model for sequences of measures I*, *Mathematical Proceedings of the Cambridge Philosophical Society* 95 (1984) 229–260.
- [MiSch] W.J. Mitchell and E. Schimmerling, *Covering without countable closure*, *Mathematical Research Letters* 2 (1995) 595–609.
- [MiSchSt] W.J. Mitchell, E. Schimmerling, and J.R. Steel, *The covering lemma up to a Woodin cardinal*, to appear in *Annals of Pure and Applied Logic*.
- [MiSt] W.J. Mitchell and J.R. Steel, *Fine structure and iteration trees*, *Lecture Notes in Logic* 3, Springer (1994).
- [N] I. Neeman, *Optimal proofs of determinacy*, *The Bulletin of Symbolic Logic* 1 (1995) 327–339.
- [Sch1] E. Schimmerling, *Combinatorial principles in the core model for one Woodin cardinal*, *Annals of Pure and Applied Logic* 74 (1995) 153–201.
- [Sch2] E. Schimmerling, $\square_\kappa^{<\omega}$ holds in $L[\vec{E}]$, submitted.
- [SchSt] E. Schimmerling and J.R. Steel, *Fine structure for tame inner models*, *The Journal of Symbolic Logic* 61 (1996) 621–639.
- [SchW] E. Schimmerling and W.H. Woodin, *The Jensen covering property*, submitted.
- [St1] J.R. Steel, *The core model iterability problem*, to appear in *Lecture Notes in Logic*.
- [St2] J.R. Steel, *Inner models with more Woodin cardinals*, *Annals of Pure and Applied Logic* 65 (1993) 185–209.
- [St3] J.R. Steel, *Core models with more Woodin cardinals*, preprint.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, IRVINE, CALIFORNIA 92697-3875, USA

E-mail address: eschimme@math.uci.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CALIFORNIA 94720, USA

E-mail address: steel@math.berkeley.edu